

Probing of nucleon mesonic structure by means of quasi-elastic knock-out processes like $p + e \rightarrow e' + \pi^+ + n$

V.G. Neudatchin, I.T. Obukhovsky*, and N.P. Yudin
Institute of Nuclear Physics, Moscow State University
Moscow 119899, Russia

February 8, 2008

Abstract

The momentum distributions $|\Psi_p^{n\pi^+}(\mathbf{k})|^2$ and $|\Psi_p^{n\rho^+}(\mathbf{k})|^2$ of pions and ρ -mesons in the nucleon are extracted from experiment. Perspectives of the quark microscopic theory of mesonic cloud are outlined.

1 Introduction

It is well known that the theoretical description of the pion photoproduction on nucleon encounters the complicated set of problems: a number and a type of used diagrams, their gauge invariance, necessity of inclusion of form factors, taking into account off-shell effects, and so on. But the situation becomes significantly simpler for virtual photons γ^* in the electroproduction process $p + \gamma^* \rightarrow n + \pi^+$ at sufficiently high values of the momentum square transfer $Q^2 = -q^2 \gtrsim 1 - 2GeV^2/c^2$. In this case the diagram with the pion pole in the t -channel (the "pion-in-flight" diagram) becomes dominant under the standard conditions of quasielastic knock-out process: $|\mathbf{q}| \gg |\mathbf{k}|$, $q_0 \gg |k_0|$, $k_0 = M_N - E_N(-\mathbf{k})$ (or $W - M_R \gg m_\pi$), where $\{k_0, \mathbf{k}\}$ is the 4-momentum of virtual pion in the nucleon and $W^2 = (p_R + p_{\pi'})^2$ is the mass of final hadron state $R + \pi$ ($R = N, \Delta, N^*$, etc.). For the first time the dominance of pion-in-flight diagram in the longitudinal part σ_L of differential cross section (integrated over the azimuthal angles) $d^3\sigma(ep \rightarrow e'n\pi^+)/dQ^2 dW^2 dt = 2\pi\Gamma(\epsilon d\sigma_L/dt + d\sigma_T/dt)$, $t \approx -\mathbf{k}^2$, was pointed in the paper [1]. Later in Ref. [2] the analysis of the momentum distribution of pions in the nucleon was given in the frame of light cone dynamics. In our works [3,4] the discussion of the knock-out process is given in the laboratory system and just in this frame we find the momentum distribution (MD) of pions (starting from experimental data [1] on σ_L). The similar situation is also for the transverse virtual photon γ_T^* (really the knockout of a vector meson with simultaneous transformation of ρ into the pion). It can provide a valuable information on the properties of vector mesons in nucleon [3]. These facts open a new possibility for the direct investigation of the meson structure of nucleon.

2 Study of the pion and ρ -meson momentum distribution in the nucleon

Our approach to the problem is very similar to the standard methods of nuclear physics, where the process of nucleon knockout has been used for a long time as a mighty tool for investigation

*The talk at International Symposium MENU2001, The George Washington Univ, 26-31 July.

of the momentum distribution of nucleons in nuclei. The cross section of quasi-elastic pion knockout from nucleon can be written in the form:

$$\frac{d\sigma_L(\gamma^*p \rightarrow n\pi^+)}{d\mathbf{k}^2} = \frac{\alpha F_\pi^2(Q^2)}{8|\mathbf{q}^*|W(W^2 - M_N^2)} \left| \frac{M(p \rightarrow n + \pi^+)}{t - m_\pi} \right|^2 \frac{\mathbf{q}^2}{Q^2} 4 \left[\omega(\mathbf{p}_{\pi'}) - q_0 \frac{\mathbf{p}_{\pi'} \cdot \mathbf{q}}{\mathbf{q}^2} \right]^2,$$

where asterisk symbols are for values written in the center of mass of the $\gamma^* + p$ collision. The "wave function" (W.F.) of a virtual pion in the nucleon is the following k-dependent factor of σ_L :

$$\left| \Psi_p^{n\pi^+}(|\mathbf{k}|) \right|^2 = \mathcal{N}^{-1} \left| \frac{M(p \rightarrow n + \pi^+)}{k_0 - \omega_\pi(|\mathbf{k}|)} \right|^2, \quad \mathcal{N}^{-1} = \frac{4\pi}{(2\pi)^3 2M_N 2E_N(|\mathbf{k}|) 2\omega_\pi(|\mathbf{k}|)},$$

which can be related to the matrix element of the pion creation operator

$$\langle n + \pi^+ | a_{\pi^+}^\dagger | p \rangle \sim \frac{M(p \rightarrow n\pi^+)}{t - m_\pi} = \frac{\mathcal{N}^{1/2}}{k_0 + \omega_\pi(|\mathbf{k}|)} \Psi_p^{n\pi^+}(|\mathbf{k}|).$$

Recall that for the standard πNN vertex the transition amplitude can be written in the form

$$\overline{M(p \rightarrow n\pi^+)}^2 = 2\mathbf{k}^2 g_{\pi NN}^2 F_{\pi NN}^2(\mathbf{k}^2), \quad F_{\pi NN}(\mathbf{k}^2) = \Lambda_\pi^2 / (\Lambda_\pi^2 + \mathbf{k}^2).$$

As the normalization of all the above factors is fixed the norm of such W.F. defines a "spectroscopic factor" (S.F.) of the $n + \pi^+$ state in the proton

$$S_p^{n\pi} = \int_0^\infty \left| \Psi_p^{n\pi^+}(k) \right|^2 k^2 dk,$$

The pion-baryon structure of nucleon can be described in terms of a set of such spectroscopic factors: $S_p^{\pi n}$, $S_p^{\pi \Delta}$, $S_p^{\pi N^*}$, etc. When the longitudinal cross section is factorizable in the form:

$$d\sigma_L(\gamma^*p \rightarrow n\pi^+)/d\mathbf{k}^2 \sim \left| \Psi_p^{n\pi^+}(|\mathbf{k}|) \right|^2 Q^2 F_\pi^2(Q^2), \quad F_\pi(Q^2) = (1 + Q^2/0.51)^{-1}$$

(e.g. in terms of the pion-in-flight mechanism) the W.F. and S.F. should be observable values in the coincidence experiments (at fixed Q^2 and M_R). Then one can define (in analogy with the standard definitions in the nuclear cluster physics) the "total number of pions in the nucleon" as a sum $S_\pi = \sum_B S_p^{\pi B}$, where $R \equiv B = N, \Delta, N^*$, etc. are all the possible virtual baryons in the nucleon. In Figs. 1 and 2 the longitudinal cross sections calculated on the basis of the above pion-in-flight mechanism (at the value of $\Lambda_\pi = 0.7$ and 1.2 GeV) are compared with the old $p(e, e'\pi^+)n$ data [1] at $Q^2 = 0.7$ and 3.3 GeV²/c². One can see that at both high and intermediate Q^2 this mechanism is in a rough agreement with the data if the cut-off parameter Λ_π is not so large (≈ 0.7 GeV/c), but new more exact data on σ_L at high Q^2 would be desirable.

The pole diagram with the ρ meson as a virtual state in the nucleon provides the main contribution to the transverse cross section $\sigma_T(\gamma^*p \rightarrow n\pi)$ at high $Q^2 \gtrsim 2 - 3$ GeV²/c² (see Fig. 3). However, the spectroscopic factor for the $n + \rho$ channel $S_p^{n\rho}$ cannot be determined from the experiment as the data are only available for too low $\mathbf{k}^2$ as compare with m_ρ^2 .

3 Pion-baryon structure of the nucleon in the 3P_0 model

Non-diagonal transitions $p \rightarrow B + \pi$ for the pion-in-flight mechanism of pion knock out can be considered on the same footing, i.e. on the basis of the field-theory vertex function, but the standard expressions

$$\left| M(p \rightarrow B_i + \pi) \right|^2 = f_{\pi NB_i}^2 \frac{4M_N M_{B_i}}{m_\pi^2} 2\mathbf{k}^2 F_{\pi NB_i}^2(\mathbf{k}^2)$$




Figure 1: Longitudinal cross section at $Q^2 = 0.7 \text{ GeV}^2/c^2$. The data are from Ref. [1]




Figure 2: Longitudinal cross section at $Q^2 = 3.3 \text{ GeV}^2/c^2$.




Figure 3: Transverse cross section at $Q^2 = 3.3 \text{ GeV}^2/c^2$.

depend on too many free parameters ($f_{\pi N B_i}$ and $F_{\pi N B_i}^2(\mathbf{k}^2)$ in each channel "i") to be informative for definite predictions of cross sections. The constituent quark model evaluations of the above free parameters would be more effective. For example, the microscopic 3P_0 model [5, 6] starts from only one free parameter, the amplitude of the vacuum $q\bar{q}$ fluctuation (normalized on the effective πqq coupling constant $f_{\pi qq} = \frac{3}{5}f_{\pi NN}$), but predict all the coupling constants $f_{\pi N B_i}$ and form factors $F_{\pi N B_i}$ of the interest starting from the standard quark-shell-model techniques [4, 7]. Parameters of the constituent quark model have already been fixed, and are not free in our approach. All the form factors $F_{\pi N B_i}$ only depend on two quark-shell-model parameters b_N and b_π (quark radii of N and π) fixed earlier. At the standard values $b_N = 0.6 \text{ fm}$ and $b_\pi = 0.3 \text{ fm}$ we have obtained the predictions for π - B_i wave functions in the nucleon (see Figs. 4 and 5), where $B_i = N, \Delta, N^* = N_{1/2-}(1535)$ and $N^{**} = N_{1/2+}(1440)$. In the used proper normalization of the W.F.'s we are able to determine the S.F.'s for these virtual states in the nucleon [7]. This values could be extracted from coincidence experiments similar to the above cited [1]. Such experiments would be more difficult because of too small predicted cross sections for $\pi + B_i$ channels, but they will be very informative for understanding the hadron structure in terms of




Figure 4: Pion momentum distribution in the nucleon for πB channels $\pi + N$ and $\pi + \Delta$.




Figure 5: Pion momentum distribution in the nucleon for πB channels $\pi + N^*$ and $\pi + N^{**}$.

quark and meson degrees of freedom.

Acknowledgments

The authors gratefully acknowledge the contribution of Dr. L.L. Sviridova to the work. The work was supported in part by the Russian Foundation for Basic Research grant No 00-02-16117.

References

- [1] B. Brauel, T. Canzler, D. Cords, et al., Z.Phys. C **3**, 101 (1979).
- [2] F. Guttner, G. Chanfray, H.P. Pirner, and B. Povh, Nucl. Phys. A**429**, 389 (1984).
- [3] V.G. Neudatchin, L.L. Sviridova, and N.P. Yudin, Yadernaya Fizika **62**, 694 (1999).
- [4] V.G. Neudatchin, I.T. Obukhovsky, and N.P. Yudin, *Relativistic Nuclear Physics and Quantum Chromodynamics*, Proc. XIV Int. Seminar on High Energy Physics Problems (Editors: A.M. Baldin and V.V. Burov), JINR, Dubna (2000), vol. 1, p. 55.
- [5] A. LeYaouanc, Ll. Oliver, O. Pène, and J.-C. Raynal, *Hadron Transitions in the Quark Model*. Gordon and Breach Sci. Publ., N.Y., London, Paris. 1988.
- [6] F. Cano, P. Gonzalez, S. Noguera, and B. Desplanques, Nucl. Phys. A**343**, 331 (1996).
- [7] I.T. Obukhovsky, The contributed paper of this issue, to be published